



Into the Future: Keeping Healthy Teeth Caries Free: Pediatric CAMBRA Protocols

FRANCISCO RAMOS-GOMEZ, DDS, MS, MPH, AND MAN-WAI NG, DDS, MPH

ABSTRACT Early childhood caries prevalence has increased significantly in children ages 2-5 years.¹ ECC disproportionately affects lower socioeconomic and minority groups, is a predictor for future decay, but is preventable and manageable.² Caries risk assessment systematically derives a patient's caries risk and is important during an infant oral health visit beginning at age 1. Information obtained through a risk assessment can guide a disease management care path tailored to an individual's age and risk to effectively treat and manage one's caries disease process.³

AUTHORS

Francisco Ramos-Gomez, DDS, MS, MPH, is a professor, University of California, Los Angeles, School of Dentistry, Section of Pediatrics, and researcher for UCSF's Center to Address Disparities in Children's Oral Health.

Man-Wai Ng, DDS, MPH, is dentist-in-chief, Children's Hospital Boston, and assistant professor, Oral and Developmental Biology, Harvard School of Dental Medicine.

A recent national health survey from 2007 found California and Texas, the two most-populated states in the United States, to rank among the lowest in children's oral health.⁴ In addition, Hispanic children are the most at risk for poor oral health, since 28.5 percent of Hispanic children compared to 19.1 percent of white children, have not seen a dentist by the age of 17. Increased awareness of the causes and consequences of ECC could help families, especially those who suffer from disparities in access to care, obtain dental care and institute preventive measures within their family practices.

Many parents and caretakers are unaware of the role they play in bacterial transmission to their child. Caregivers pass organisms and bacteria to infants

orally through close contact (vertical transmission).⁵ Furthermore, women from vulnerable underserved communities, and some of their providers, fail to recognize the value of good oral health and relevant importance of regular dental visits and care during pregnancy.⁶

It is critical that oral health providers, whether at academic centers, in private practice, or at safety net sites (such as health centers and hospital clinics) embrace risk assessment and disease management in addressing ECC. In its Life Course Model 2010 concept paper, "A conceptual framework that helps explain health and disease patterns particularly health disparities — across populations and over time," the U.S. Department of Health and Human Services Maternal and Child Health Bureau (MCHB) posits,

“that interventions that reduce risks and increase protective factors can change the health trajectory of individuals and populations.” It further suggested “the need to: refocus resources and strategies for a greater emphasis on early (“upstream”) determinants of health; incorporate earlier detection of risks coupled with earlier intervention; and promote protective factors while reducing risk factors at the individual, family, and community levels.”^{3,7}

Caries management by risk assessment (CAMBRA) is designed for use with newborns to 5-year-old children. It is easy to use and offers an approach to disease prevention management that integrates risk assessment of childhood caries as an integral component of a comprehensive oral health visit.

Caries Management by Risk Assessment

CAMBRA assists providers to systematically:

- Assess each child and his caregiver’s caries risk in an individualized manner;
- Tailor a specific preventive therapeutic management plan or “care path”;
- Customize a restorative plan in conjunction with preventive care; and
- Plan a timely, specific, and appropriate periodicity schedule based on the child’s caries risk.

To effectively prevent and management the disease of caries, care should begin early, ideally during an age 1 comprehensive oral exam visit. There are six basic steps in the infant oral care visit. A caries risk assessment is the first of these six critical steps, giving the provider more information to help them consider the risk and health status of each patient before beginning the exam.³

Caries risk assessment provides information pertaining to three specific overarching domains:

Risk and/or biological factors such as continual bottle use, sleeping with a bottle, frequency and types of snacks, a child taking any medications as well as other risk factors;

Protective factors such as the use of fluoridated tap water, use of fluoridated toothpaste, or the use of xylitol on a continuous basis; and

Clinical findings such as the presence of early demineralized enamel surface, cavitated lesions, plaque, lack of salivary flow, etc. (informa-

CARIES RISK
assessment can
be easily and
efficiently performed
by dental and
medical providers.

tion to be obtained from Step 4).

Through a short and brief interview with the caregiver, information is gathered to assess the child’s risk of caries development and disease progression as low, moderate, or high. For example, a child may be at high risk if the child goes to bed or has a constant exposure with a bottle containing liquids with natural or artificial sugar, or snacks throughout the day, etc. Protective factors include brushing with a smear of fluoridated toothpaste at least once daily, especially before bed at night or drinking fluoridated tap water regularly.

Three findings are always associated with a high caries risk. These are: 1) new carious lesions in the primary caregiver within the past 12 months; 2) prior caries and/or restorations in the child; and 3)

white spot lesions, decalcification enamel defects or other obvious decay in the child (**TABLE 1**). The information obtained from a caries risk assessment allows the care provider to formulate a caries risk profile for the child, an essential first step to determining the prevention and treatment plan, as well as the periodicity of patient follow-up/recall (one month, three months, six months, or one year).³ Caries risk assessment can be easily and efficiently performed by dental and medical providers.

Step 2 consists of proper positioning of the infant. The knee-to-knee position allows an effective and efficient visualization of the child’s oral cavity and dentition. The child is laying supine, with his head resting in the care provider’s lap. This position allows the child to see his parent and the parent to see what the care provider sees.

Step 3 involves a toothbrush prophylaxis, which is effective in the removal of plaque on most teeth. Using the tell-show-do technique, the care provider can demonstrate the proper technique for brushing the child’s teeth. Step 4 is the clinical examination and Step 5 is the application of fluoride varnish, which is to prevent tooth decay. Fluoride varnish is to be applied every three to six months depending on the caries risk of the child.^{3,8,9}

Step 6 involves the care provider working with the parent to determine a mutually agreed upon set of self-management goals appropriate for the family (**FIGURE 1**). The care provider is to first transmit a summary of the findings obtained from the caries risk assessment and clinical examination, and explain to the parent the causes of the caries process. Together with the parents, one or two self-management goals are to be identified for the family to work on to reduce the child’s risk factors and enhance his protective factors.³ Parents should be

TABLE 1

CAMBRA — Caries Risk Assessment Form for Age 0 to 5 Years

Patient Name: _____ ID# _____ Age: _____ Date: _____

Assessment Date: _____ Please circle: BASELINE, three-month follow-up or six-month follow-up

	1	2	3	
NOTE: Any one Yes in Column 1 signifies likely "High Risk" and an indication for bacteria tests	Yes =CIRCLE	Yes =CIRCLE	Yes =CIRCLE	Comments:
1. Risk Factors (Biological Predisposing Factors)				
(a) Mother or primary caregiver has had active dental decay in the past 12 months*	Yes			
(b) Bottle with fluid <u>other</u> than water, plain milk and/or plain formula		Yes		Type of fluid:
(c) Continual bottle use		Yes		
(d) Child sleeps with a bottle, or nurses on demand		Yes		
(e) Frequent (>3 times/day) between-meal snacks of sugars/cooked starch/sugared beverages		Yes		#times/day:
(F) Saliva-reducing factors are present, including: 1. medications (e.g., some for asthma [albuterol] or hyperactivity) 2. medical (cancer treatment) or genetic factors		Yes		
(g) Child has developmental problems/CSHCN (child with special health care needs)		Yes		
(h) Caregiver has low health literacy, is a WIC participant and/or child participates in Free Lunch Program and/or Early Head Start		Yes		
2. Protective Factors				
(a) Child lives in a fluoridated community or takes fluoride supplements by slowly dissolving or as chewable tablets (note resident ZIP code)			Yes	
(b) Child drinks fluoridated water (e.g., use of tap water)			Yes	
(c) Teeth brushed with fluoridated toothpaste (pea-size) at least once daily			Yes	
(d) Teeth brushed with fluoride toothpaste (pea-size) at least 2x daily			Yes	
(e) Fluoride varnish in last six months			Yes	
(f) Mother/caregiver chews/dissolves xylitol chewing gum/lozenges 2-4x daily			Yes	
3. Disease Indicators/Risk Factors – Clinical Examination of Child				
(a) Obvious white spots, decalcifications enamel defects or obvious decay present on the child's teeth*	Yes			
(b) Restorations present (past caries experience for the child)*	Yes			
(c) Plaque is obvious on the teeth and/or gums bleed easily		Yes		
(d) Visually inadequate saliva flow		Yes		
(e) New remineralization since last exam (List teeth):				

Child's Overall Caries Risk* (circle): High Moderate Low

Child: Bacteria/Saliva Test Results: MS: LB: Flow Rate: ml/min: Date:

Caregiver: Bacteria/Saliva Test Results: MS: LB: Flow Rate: ml/min: Date:

Self-management goals:

- 1) _____
- 2) _____



**VISUALIZE
CARIES BALANCE**


*Assessment based on provider's judgment of balance between risk factors/disease indicators and protective factors.

Doctor signature/#: _____ Date: _____

Self-Management Goals for Parent/Caregiver

Patient Name _____

DOB _____



Regular dental visits for child



Family receives dental treatment



Healthy snacks



Brush with fluoride toothpaste at least 2 times daily



No soda



Less or no juice



Wean off bottle (no bottles for sleeping)



Only water or milk in sippy cups



Drink tap water



Less or no junk food and candy



Use xylitol spray, gel or dissolving tablets

IMPORTANT: The last thing that touches your child's teeth before bedtime is the toothbrush with fluoride toothpaste.

Self-management goals 1) _____
 2) _____

On a scale of 1-10, how confident are you that you can accomplish the goals? 1 2 3 4 5 6 7 8 9 10

Signature _____ Date _____

Practitioner signature _____ Date _____

FIGURE 1. Self-management goals.

encouraged to adopt a healthy oral lifestyle for their child, assist their children from an early age to maintain a proper diet and oral hygiene, and establish and maintain a dental home for their child.³ When self-management goals are revisited during recall visits, caregivers and their children can receive positive reinforcement as they see how meeting their goals can improve their child's caries risk and result in better oral health outcomes.

Risk-Based Care Paths

Once a child's caries risk has been determined, the care provider, in partnership with the child's caregiver, can determine a multifaceted care path appropriate for the family, based on the child's age and individualized needs. Research supports the use of fluoride varnish in combination with improved diet and oral hygiene counseling, and families should be encouraged to drink fluoridated tap water and/or use fluoridated toothpaste, which are very important aspects and measures of preventive care.^{8,9} However, a care path or decision tree can aid the provider in determining a specific combination of diagnostic, preventive and restorative procedures, and the periodicity of these recommended measures that are appropriate for the child and his family to improve and/or stabilize a child's caries risk profile.

Care paths are expected to be dynamic and change over time as the effectiveness of new as well as current protocols are validated by scientific evidence. **TABLES 2 AND 3** are examples of the care paths at this time recommended for children ages 0-2 and ages 3-5, respectively. Newer products, such as remineralizing gels, or new uses for existing products such as glass ionomer to be used as sealants, are now available and can be considered viable treatment options. As evidence continues to evolve, and new studies advocate for the use of

agents, such as probiotics and topical applications of providone-iodine to reduce high oral bacteria levels, these modalities will be added to the care path. While the evidence for the effectiveness of preventive and treatment modalities continues to grow, standardized and widely accepted protocols are limited. Therefore, each practitioner should use careful consideration based on emerging available evidence and one's own experience when considering when and how to introduce use of newer modalities into their patients' care paths.

FAMILIES SHOULD BE encouraged to drink fluoridated tap water and/or use fluoridated toothpaste, which are very important aspects and measures of preventive care.

Quality Improvement — Potential for Transforming Oral Health Care Delivery and Improving Outcomes

Evidence supports the effectiveness of dietary control, fluoride use, and other modalities such as xylitol, in preventing and controlling dental caries, a chronic infectious disease caused by acids produced by oral bacteria metabolizing fermentable carbohydrates.¹⁰ Since the risk for caries development and caries activity differs among individuals and may change in each individual over time, caries risk assessment performed initially, and periodically thereafter, allows for a determination of a patient's relative risk, from which an adoption of an evidence-based prevention plan that can be customized.

Disease management of caries (e.g., CAMBRA) is modeled after the medical

management of chronic conditions in which the patient (or child's family) is engaged in day-to-day health behavior modifications that address disease etiology. It requires parent/family engagement in dietary control and applications of fluorides and other preventive modalities. The progress in stabilizing and reducing the risk for the caries process is monitored using caries risk assessment tools.

Despite the evidence supporting the effectiveness of bio-behavioral approaches and interventions in preventing and controlling caries, including caries risk management, to improve patient outcomes, disease management of caries has not been widely implemented in clinical dental practice. It is known that barriers exist in our current oral care delivery systems, which do not easily permit effective implementation of disease management of caries in clinical practice. These barriers include a provider's lack of knowledge, skills and comfort; a parent's knowledge, preferences and expectations; reimbursement favoring surgical management of caries; and coordination and follow-up. While reimbursement methodologies need to be altered from the current "one-size-fits all" to include coverage of caries risk assessment and more frequent and intensive preventive services for higher-risk individuals, quality improvement (QI) can foster and accelerate the adoption of changes needed to redesign current care delivery systems. QI methods also have allowed for the testing of system changes in order to produce better system performance and improved outcomes for patients and populations.

QI is defined by Batalden as the combined and unceasing efforts of everyone — health care professionals, patients and their families, researchers, payers, planners and educators — to make changes that will lead to better patient outcomes (health), better system performance (care) and better pro-

TABLE 2

Example of a Caries Management Protocol for 0–2 Year Olds

Risk Category Ages 0 to 2	Diagnostic			Fluoride
	Periodic Oral Exams	Radiographs*	Saliva Test	
Low	Annual	Posterior bitewings at 12-24 month intervals if proximal surfaces cannot be examined visually or with a probe	Optional baseline	In office: No Home: Brush 2x day w/ smear of F toothpaste
Moderate	Every 6 months	Posterior bitewings at 6-12 month intervals if proximal surfaces cannot be examined visually or with a probe	Suggested	In office: F varnish initial visit & recalls Home: Brush 2x day w/smear of F toothpaste Caregiver: OTC sodium fluoride treatment rinses
Moderate; noncompliant	Every 3-6 months	Posterior bitewings at 6-12 month intervals if proximal surfaces cannot be examined visually or with a probe	Recommended	In office: F varnish initial visit & recalls Home: Brush 2x day w/smear of F toothpaste then apply a smear of calcium-phosphate paste left on at bedtime Caregiver: OTC sodium fluoride treatment rinses
High	Every 3 months	Anterior (#2 occlusal film) and posterior bitewings at 6-12 month intervals if proximal surfaces cannot be examined visually or with a probe	Recommended	In office: F varnish initial visit & recalls Home: Brush 2x day w/smear of F toothpaste then apply a smear of calcium-phosphate paste left on at bedtime Caregiver: OTC sodium fluoride treatment rinses
High; noncompliant	Every 1-3 months	Anterior (#2 occlusal film) and posterior bitewings at 6-12 month intervals if proximal surfaces cannot be examined visually or with a probe	Recommended	In office: F varnish initial visit & recalls Home: Brush 2x day w/smear of F toothpaste then apply a smear of calcium-phosphate paste left on at bedtime Caregiver: OTC sodium fluoride treatment rinses
Extreme	Every 1-3 months	Anterior (#2 occlusal film) and posterior bitewings at 6-12 month intervals if proximal surfaces cannot be examined visually or with a probe	Recommended	In office: F varnish initial visit & recalls Home: Brush 2x day w/smear of F toothpaste then apply a smear of calcium-phosphate paste left on at bedtime Caregiver: OTC sodium fluoride treatment rinses

Preventive Intervention						Restoration**
Xylitol	Sealants	Antibacterials/ Probiotics	Anticipatory Guidance/ Counseling	Self- Management Goals	White Spot / Precavitated Lesions	Existing Lesions
Not Required	No	No	Yes	No	n/a	n/a
Child: Xylitol wipes 3-4x daily Caregiver: 2 sticks of gum or 2 mints 4x day, total 6-10 grams a day	Glass ionomer- based materials recommended on deep pits and fissures	No	Yes	No	Treat w/ fluoride products as indicated to promote remineralization	n/a
Child: Xylitol wipes 3-4x daily Caregiver: 2 sticks of gum or 2 mints 4x day, total 6-10 grams a day	Glass ionomer- based materials recommended on deep pits and fissures	Recommend CHX for caregiver/ Recommend probiotics	Yes	Yes	Treat w/ fluoride products as indicated to promote remineralization	n/a
Child: Xylitol wipes 3-4x daily Caregiver: 2 sticks of gum or 2 mints 4x day, total 6-10 grams a day	Glass ionomer- based materials recommended on deep pits and fissures	Recommend CHX for caregiver/ Recommend probiotics	Yes	Yes	Treat w/ fluoride products as indicated to promote remineralization	ITR (interim therapeutic restorations) with glass ionomer-based materials or conventional restor- ative treatment as patient cooperation and family circumstances allow
Child: Xylitol wipes 3-4x daily Caregiver: 2 sticks of gum or 2 mints 4x day, total 6-10 grams a day	Glass ionomer- based materials recommended on deep pits and fissures	Recommend CHX for caregiver/ Recommend probiotics	Yes	Yes	Treat w/ fluoride products as indicated to promote remineralization	ITR (interim therapeutic restorations) with glass ionomer-based materials or conventional restor- ative treatment as patient cooperation and family circumstances allow
Child: Xylitol wipes 3-4x daily Caregiver: 2 sticks of gum or 2 mints 4x day, total 6-10 grams a day	Glass ionomer- based materials recommended on deep pits and fissures	Recommend CHX for caregiver/ Recommend probiotics	Yes	Yes	Treat w/ fluoride products as indicated to promote remineralization	ITR (interim therapeutic restorations) with glass ionomer-based materials or conventional restor- ative treatment as patient cooperation and family circumstances allow

TABLE 3

Example of a Caries Management Protocol for 3-5-Year-Olds

Risk Category —Ages 3 to 5	Diagnostic			Fluoride
	Periodic Oral Exams	Radiographs*	Saliva Test	
Low	Annual	Posterior bitewings at 12-24 month intervals if proximal surfaces cannot be examined visually or with a probe	Optional baseline	In office: No Home: Brush 2x day w/ pea-size amount of F toothpaste
Moderate	Every 6 months	Posterior bitewings at 6-12 month intervals if proximal surfaces cannot be examined visually or with a probe	Suggested	In office: F Varnish initial visit & recalls Home: Brush 2x day w/ pea-size amount of F toothpaste Caregiver: OTC sodium fluoride treatment rinses
Moderate; noncompliant	Every 3-6 months	Posterior bitewings at 6-12 month intervals if proximal surfaces cannot be examined visually or with a probe	Recommended	In office: F varnish initial visit & recalls Home: Brush 2x day w/ pea-size amount of F toothpaste, then apply pea-size amount of calcium-phosphate paste left on at bedtime Caregiver: OTC sodium fluoride treatment rinses
High	Every 3 months	Anterior (#2 occlusal film) and posterior bitewings at 6-12 month intervals if proximal surfaces cannot be examined visually or with a probe	Recommended	In office: F varnish initial visit & recalls Home: Brush 2x day w/ pea-size amount of F toothpaste, then apply pea-size amount of calcium-phosphate paste left on at bedtime Caregiver: OTC sodium fluoride treatment rinses
High; noncompliant	Every 1-3 months	Anterior (#2 occlusal film) and posterior bitewings at 6-12 month intervals if proximal surfaces cannot be examined visually or with a probe	Recommended	In office: F varnish initial visit & recalls Home: Brush 2x day w/ pea-size amount of F toothpaste, then apply pea-size amount of calcium-phosphate paste left on at bedtime Caregiver: OTC sodium fluoride treatment rinses
Extreme	Every 1-3 months	Anterior (#2 occlusal film) and posterior bitewings at 6-12 month intervals if proximal surfaces cannot be examined visually or with a probe	Recommended	In office: F varnish initial visit & recalls Home: Brush 2x day w/ pea-size amount of F toothpaste, then apply pea-size amount of calcium-phosphate paste left on at bedtime Caregiver: OTC sodium fluoride treatment rinses

Preventive Intervention						Restoration**
Xylitol &/or Baking Soda	Sealants	Antibacterials/ Probiotics	Anticipatory Guidance/ Counseling	Self-Management Goals	White Spot / Precavitated Lesions	Existing Lesions
Not Required	No	No	Yes	No	n/a	n/a
Child: Xylitol wipes 3-4x daily; products to substitute for sweet treats or when unable to brush Caregiver: 2 sticks of gum or 2 mints 4x day, total 6-10 grams a day	Glass ionomer-based materials recommended on deep pits and fissures	No	Yes	No	Treat w/fluoride products as indicated to promote remineralization	n/a
Child: Xylitol wipes 3-4x daily; products to substitute for sweet treats or when unable to brush Caregiver: 2 sticks of gum or 2 mints 4x day, total 6-10 grams a day	Glass ionomer-based materials recommended on deep pits and fissures	Recommend CHX for caregiver/ Recommend probiotics	Yes	Yes	Treat w/fluoride products as indicated to promote remineralization	n/a
Child: Xylitol wipes 3-4x daily; products to substitute for sweet treats or when unable to brush Caregiver: 2 sticks of gum or 2 mints 4x day, total 6-10 grams a day	Glass ionomer-based materials recommended on deep pits and fissures	Recommend CHX for caregiver/ Recommend probiotics	Yes	Yes	Treat w/fluoride products as indicated to promote remineralization	ITR (interim therapeutic restorations) with glass ionomer-based materials or conventional restorative treatment as patient cooperation and family circumstances allow
Child: Xylitol wipes/ products to substitute for sweet treats or when unable to brush Caregiver: 2 sticks of gum or 2 mints 4x day, total 6-10 grams a day	Glass ionomer-based materials recommended on deep pits and fissures	Recommend CHX for caregiver/ Recommend probiotics	Yes	Yes	Treat w/fluoride products as indicated to promote remineralization	ITR (interim therapeutic restorations) with glass ionomer-based materials or conventional restorative treatment as patient cooperation and family circumstances allow
Child: Xylitol wipes 3-4x daily; products to substitute for sweet treats or when unable to brush Caregiver: 2 sticks of gum or 2 mints 4x day, total 6-10 grams a day	Glass ionomer-based materials recommended on deep pits and fissures	Recommend CHX for caregiver/ Recommend probiotics	Yes	Yes	Treat w/fluoride products as indicated to promote remineralization	ITR (interim therapeutic restorations) with glass ionomer-based materials or conventional restorative treatment as patient cooperation and family circumstances allow

fessional development (learning).¹¹ Quality improvement necessarily involves making changes that systematically incorporate evidence-based knowledge. It functions at the system level by which care delivery takes place, the physical and information level along with the complex social structures — i.e., health care professionals. In recent years, hospitals and medical health care systems have increasingly been using QI to effectively enhance patient safety, improve quality of care, and management of chronic disease and preventive care.

Although QI is not yet familiar territory to dentistry, it offers the potential to transform oral health care delivery in order to provide better oral health care, improve oral health outcomes, and to reduce costs of treatment of caries. A demonstration project funded by the DentaQuest Institute was implemented at two hospital-based dental practices that care for a disproportionate number of young children with ECC to test the feasibility of applying a disease management approach for ECC. Applying the principles of QI, it focused on making changes within the practices' oral health care delivery systems needed to implement an evidence-based disease management approach for ECC. Thirty months of results from the demonstration project have shown that a disease management approach to address ECC can be implemented into practice and has the potential to deliver better care, improve clinical outcomes, and reduce the overall cost of care.¹²

Ultimately, in order for the caries risk assessment to be successfully implemented as a universal model for quality improvement, the public and private systems of care must enact an equitable financial reimbursement model for these preventive treatments and its intervention care paths based on a medical model based on age and risk that is comparable and/or equitable to current surgical care compensation. Fur-

thermore, providers and insurers alike must embrace the dental ethics perspective of early disease prevention and early intervention that can benefit their future patients' short- and long-term oral health outcomes.

Summary

While caries risk assessment was first endorsed by the American Dental Association, the American Academy of Pediatric Dentistry, and the American Association of Public Health Dentistry, its use by nondenial professionals has become more

A DISEASE MANAGEMENT approach to address ECC can be implemented into practice and has the potential to deliver better care, improve clinical outcomes, and reduce the overall cost of care.

widespread.¹³⁻¹⁵ The American Academy of Pediatrics now recommends the use of a risk assessment protocol during well-child visits to all its providers.¹⁶ Community-based organizations, principally those working with underserved and vulnerable populations such as Early Head Start and Women, Infants and Children have found CAMBRA an essential tool as part of their comprehensive infant oral care programs.

CAMBRA's easy-to-use organized format of disease indicators, risk and protective factors, clinical findings, and self-management goals helps to facilitate oral health education, deepens the appreciation of oral health information, and increases the understanding of how individual behaviors can affect caries development and progression. By

embracing the concepts of caries risk assessment, early intervention and the establishment of a dental home, care providers could reduce their patients' risk of early childhood caries and improve children's oral and overall health.

The adoption of CAMBRA has not been universal in the dental community, which remains primarily focused on restorative treatment rather than on prevention and management of the disease. Since restorative and surgical care by itself does not address disease etiology, patients and caregivers must have the opportunity to increase their own oral health literacy, understand the causes and consequences of poor oral health and the value of "maintaining healthy teeth." As families gain an understanding of the value of preventive care and managing their oral disease risk, they will expect to partner with their oral health provider on plans for preventive care, monitoring, early intervention and treatments, as they have come to demand with their physicians. As QI becomes more familiar to dentistry, dental practices will be able to use QI methods to redesign their systems of care and train themselves and their staff to deliver customized risk-based prevention and disease management to patients. In doing so, the dental profession will have the potential to improve patient oral health outcomes and result in a reduction in cost of care.

It is important to recognize that, in order for a successful paradigm shift to disease prevention to occur full scale, dental insurance benefits must support risk-based disease management by reimbursing for risk assessments, monitoring and regularly assessing, and more frequent preventive care treatments for high caries risk patients. Despite these challenges, many practices and clinics have successfully implemented CAMBRA. Two other articles will be presented in the *Journal* as helpful guidelines to CAMBRA adoption and incorporation into practice. ■■■■